

PREVENTION OF DIABETES**FIELD OF THE INVENTION**

5 The present invention relates to the use of an insulin-like polypeptide without hypoglycaemic effect for delaying or preventing the onset of diabetes or for ameliorating an early stage thereof.

10

BACKGROUND OF THE INVENTION

Autoimmune diseases are characterized by an abnormal immune response (involving either immune system cells or
15 antibodies) directed against normal autologous (self) tissues. Autoimmune diseases afflict huge numbers of individuals throughout the world.

A normal immune system has the capacity to identify and
20 destroy a large variety of foreign invader organisms such as bacteria and viruses. Remarkably, a normal immune system can readily distinguish foreign substances from self, and thereby is able to react vigorously against potentially pathogenic entities from the environment without harming the
25 host's own cells.

The immune system's non-reactivity to self is termed immunological tolerance. In pathological situations, immunological tolerance to a wide variety of self substances is broken,
30 resulting in an autoimmune response. If of an appropriate nature and of sufficient severity and duration, the anti-self response will result in an autoimmune disease. Cellular immune mechanisms are believed to be primarily involved in insulin dependent diabetes (IDD).

35

Whereas susceptibility to autoimmune diseases may be inherited through the defective actions of multiple genes, indirect evidence suggests that an interaction with a

foreign substance from the environment may also be necessary to induce the pathogenic process that results in disease. One explanation for this is that immunization with the foreign inductive chemical induces a cross-reactive response to self through molecular mimicry or chemical similarity. However, once the autoimmune process has been initiated, other secondary immunizing events involving other self antigens typically occur through the release of intracellular constituents in forms not normally encountered by the immune system. Targeted organs thus become damaged through the combination of all of these events, which leads to the appearance of a clinically recognized disorder only when the disease process has progressed to ablate large numbers of tissue cells so targeted.

A number of strategies have been used or proposed to suppress autoimmune diseases, most notably drugs, such as cyclophosphamide, cyclosporin A, methotrexate, and azathioprine. Steroid compounds, such as prednisone and methylprednisolone, are also employed in many instances. These drugs have limited long term efficacy against both cell- and antibody-mediated autoimmune diseases. Use of drugs is limited by virtue of their toxic side effects which include universal immunosuppression. Prolonged treatment with these drugs inhibits the normal protective immune response to pathogenic microorganisms, thereby increasing the risk of infections. A further drawback is that immune-mediated elimination of aberrant cells is impaired and there is, thus, an increased risk that malignancies will develop in patients receiving prolonged global immunosuppression.

Because the subject invention concerns insulin dependent (Type I) diabetes, a detailed background of diabetes is provided below.

Insulin dependent diabetes. Diabetes mellitus comprises a group of diseases that result in elevation of the blood glucose level because of relative or absolute deficiency in

the pancreatic hormone insulin. Insulin is secreted into the blood when food is ingested and primarily directs absorbed nutrients into body stores. Diabetes is a major public health problem affecting at least 5 million and as many as 10 million Americans. The prevalence of the most severe form of IDD is 1 in 300 in the United States.

Chronic elevation of the blood glucose level is the most obvious metabolic effect in diabetes and is associated with progressive damage to blood vessels. This may lead to heart attack, stroke, blindness, peripheral nerve dysfunction, and kidney failure. The frequency and severity of diabetes-related complications are greatest in the insulin dependent form of the disease, in which an immunological destruction of the insulin secreting pancreatic beta cells occurs. The high rate of irreversible complications in IDD occurs despite the availability of insulin replacement through injections given 1-4 times daily.

Insulin and other pancreatic hormones are well known and characterized. See, for example, Steiner et al. (1989) "Chemistry and Biosynthesis of Pancreatic Protein Hormones," in Endocrinology; DeGroot et al., EDs., W.B. Saunders Company, p. 1263-1289. As described in Steiner et al., the amino acid sequence of insulin is highly conserved across a number of species, including human, monkey, swine, and ox. IDD has proved itself to be predictable both in unaffected relatives of patients with IDD, as well as in persons from the general population. A predisposition to develop clinical diabetes can be detected through several different tests. For example, genetic susceptibility to diabetes has become increasingly definable through the use of molecular biological means, usually from DNA samples obtained from peripheral blood. One major gene involved in the inherited susceptibility to IDD is that located at the HLA-DQ locus. It is currently possible to identify risks varying from essentially none to those as high as 70 fold above those without the genotype. In families a genetic risk as high as

1 in 4 can be estimated for unaffected siblings just through identification of HLA haplotypes shared with the affected proband.

- 5 Persons who have just developed IDD or are in process of developing IDD have a number of disease-specific autoantibodies in their blood. Such autoantibodies include those to islet cell antigens (ICA), to beta cell specific proteins of 64 kDa, which are now believed to be the lower
10 molecular isoform of glutamic acid decarboxylase (GAD₆₅), to native insulin and proinsulin, and to a number of more minor determinants such as carboxypeptidase-H and heat shock proteins belonging to the hsp-60 family.
- 15 Insulin autoantibodies (IAA) are observed in untreated, newly diagnosed IDD patients (Palmer et al. (1983) Science **222**, 1337-1339) as well as in apparently unaffected relatives of diabetic probands. Whereas autoimmunity to insulin could directly cause beta-cell damage, could
20 interfere with the action of endogenous insulin, or could have both effects, some investigators suggested that IAA reflect the rate of islet cell destruction and thus act merely as reporters of aggressive islet directed autoimmunity (Ziegler et al. (1989) Diabetes **38**, 1320-1325;
25 Vardi (1988) Diabetes Care **11**, 736-739).

- The spontaneously diabetic non-obese diabetic (NOD) mouse and the BB rat are useful animal models for human IDD. Analysis of these animals provides important insights into
30 the sequence of pathogenic events, and leads to an understanding of the nature of the autoimmunological process. Previous studies from several laboratories have demonstrated that an extended prophylactic course of daily, subcutaneous injections of high doses of insulin protected
35 NOD mice and BB rat from both hyperglycaemia and islet infiltration by mononuclear leucocytes (insulinitis) (Atkinson et al. (1990) Diabetes **39**, 933-937, Gotfredsen et al. (1985) Diabetologia **28**, 933-935).

In addition, prophylactic treatment with high doses of insulin has been reported to prevent diabetes in NOD mice and in BB rats adoptively transferred with spleen lymphocytes from acutely diabetic animals to their non-diabetic counterparts (Thivolet et al. (1991) *Diabetologia* **34**, 314-319).

Such treatment may relieve the pancreatic beta-cells of metabolic demands and thus induce a state of "beta-cell rest". This quiescent state may be associated with diminished expression of many islet factors, including those that may serve as potential autoantigens at the cell surface (Aaen et al. (1990) *Diabetes* **39**, 697-701; Kämpe et al. (1989) *Diabetes* **38**, 1326-1328). Non-specific immunostimulation caused by cytokine (Jacob et al. (1990) *Proc. Natl. Acad. Sci. USA* **87**, 968-972) or adjuvant (Sadelain et al. (1990) *Diabetes* **39**, 583-589) treatments, or environmental microbes have been implicated in other protocols of IDD prevention (Like et al. (1990) *Diabetes* **40**, 259-262). A better understanding of the pathogenic role of insulin autoimmunity in IDD is clearly required.

There have been reports of efforts to induce antigen-specific immunoregulation to ameliorate autoimmune diseases (Steinman, L. (1990) *Mol. Biol. Med.* **7**, 333-339). For example, various methods have been employed to induce antigen-specific suppression of experimental allergic encephalomyelitis (EAE) (PCT publication WO 91/15225). Recently, several novel immunological approaches have been explored to autoimmune diseases such as EAE in mice and rats and lupus nephritis in MRL/lpr mice. Many have been directed toward blocking the function of the effector CD4⁺ T cell which has been shown to exhibit V_β isotype restriction in EAE. These approaches have included the use of anti-TCR antibodies (Archa-Orbea et al., *supra*), synthetic TCR peptides (Offner et al. (1991) *Science* **251**, 430-432) and superantigen treatment (Kim et al. (1991) *J. Exp. Med.* **174**,

1431-1437). The tolerogenic effects of enteral or pulmonary administration of antigens have also been described (Silverman et al. (1983) J. Immunol. **131**, 2651-2661; Peng et al. (1990) Clin. Exp. Immunol. **81**, 510-515; Michael (1989) Immune Invest. **18**, 1049-1054; Kitamura et al. (1987) J. Immunol. **139**, 3251-3259; Michalek et al. (1982) J. Immunol. **128**, 1992-1998; Mowat et al. (1986) Immunol. **58**, 677-683; PCT publications WO 91/12816, WO 91/01333; WO 92/06704). Nagler-Anderson et al. (1986) Proc. Natl. Acad. Sci. USA **83**, 7443-7446 describe the suppression of arthritis by oral administration of soluble type II collagen in a collagen-induced arthritis mouse model. Deficiencies in this ability have been reported in several experimental mouse models of autoimmune diseases (Gesualdo et al. (1990) J. Immunol. **145**, 3684-3691; Miller et al. (1984) Clin. Immunol. Immunopathol. **31**, 231-240).

Zhang et al. have in NOD mice observed a reduced frequency of diabetes in response to very high doses of oral insulin (Zhang et al. (1991) Proc. Natl. Acad. Sci. USA **88**, 10252-10256) and further demonstrated in co-transfer studies that splenocytes from insulin-fed NOD mice prevented the adoptive transfer of diabetes by splenocytes from untreated diabetic mice to irradiated recipients.

In contrast, oral administration of insulin to spontaneously or induced diabetes-prone BB rats was reported not to offer any protection against diabetes (Mordes et al., 1995, The New York Academy of Science, "Oral Tolerance: Mechanism and Application").

Muir et al. (PCT publication WO 94/23737) have reported that, in NOD mice, administration of human insulin or its B-chain together with a powerful immune-stimulating adjuvant provides a small but significant protection against diabetes. This protection can be adoptively transferred to irradiated NOD mice co-infused with diabetogenic spleen cells and is associated with a reduced expression of gamma-

interferon in the islet infiltrating lesion.

Also in the NOD mouse, nasal administration of insulin or a fragment of its B-chain has been reported to protect against diabetes adoptively transferred to non-diabetic NOD mice with a diabetogenic NOD T cell clone reactive against an amino acid sequence harboured in the insulin B-chain fragment referred to above or against spontaneous diabetes in NOD mice (Wegman et al. (1995) The New York Academy of Science, "Oral Tolerance: Mechanism and Application").

In yet another example of the use of insulin in the protection against diabetes, it has been reported that a single intravenous injection of insulin into NOD mice is followed by a reduced incidence of diabetes compared to non-treated animals.

The above examples of using insulin for preventing the onset of clinical diabetes have all been carried out in animal models of IDD, i.e. the NOD mouse and BB rat, but there is an increasing evidence that the same phenomena occur in the case of human IDD. Thus, fewer individuals, at high risk of developing IDD based on family history and immune markers, that were treated in the non-diabetic state with daily s.c. injections of insulin developed IDD than untreated individuals at a comparable high risk. Currently, large clinical studies in the US and in Europe are undertaken to formally prove whether prophylactic insulin therapy in non-diabetic, at-risk individuals can protect against subsequent development of IDD.

The mechanism whereby insulin exerts its effect in protecting against diabetes are not known exactly, but two issues are common to the above mentioned examples of the prophylactic use of insulin. One is that in most cases hormonally active insulin has been used in doses sufficient to cause constant or transient hypoglycaemia. In fact, the doses used in the animals studies have often led to high frequencies of deaths due to hypoglycaemia and the doses

used in humans are of a magnitude where hypoglycaemic symptoms may frequently be expected.

The other common issue is the fact that a non-hormonal, immunologic protective mechanism of insulin, e.g., nasal administration of insulin B-chain fragment, relies on the use of a genetically homogenous animal strain. The immune response to an antigenic stimulus, whether of immunizing or tolerogenic nature, occurs via the presentation of a peptide fragment of the protein antigen in question to T cells. The peptide is presented on a cellular receptor, the MHC (major histocompatibility) molecules. The MHC molecules are extremely polymorphic, thus two classes of molecules exist, class I and class II and within each class several loci exist, e.g., the HLA class I (A, B and C loci) and HLA class II (DR, DQ and DP) in humans, and within each locus a large number of polymorphic alleles exist. Different MHC molecules present different peptides to T cells, even from the same antigen. Consequently, to obtain a T cell response, be it immunogenic or tolerogenic, to a given antigen, e.g., insulin in an outbred, MHC polymorphic population like humans, administration of the entire antigen, or at least the major part of it is likely to be required to obtain the desired response in all or most individuals treated.

According to the present invention, an insulin-like polypeptide without hypoglycaemic effect is used to prevent against the development of diabetes. The insulin is rendered hormonally inactive by discrete, specified deletions or amino acid substitutions which leaves the insulin minimally changed or as a fragment of the endogenous molecule. The present invention makes possible the treatment of clinical healthy individuals, albeit at a certain risk of developing a serious disease, with a disease specific antigen, insulin, without exposing the individual to the potential serious adverse effects, hypoglycaemia and mitogenicity secondary to the actions of hormonally active insulin. In an example of the present invention, daily sc injections with human

insulin with a single amino acid substitution in position B25 that renders the molecule less than 0.1% hormonally active, in NOD mice protects as well against diabetes as equimolar doses of human insulin does, but without the adverse effects of insulin, i.e., approx. 50% deaths due to hypoglycaemia.

SUMMARY OF THE INVENTION

10

In one aspect, the present invention concerns insulin molecules rendered hormonally inactive by specific deletions and/or amino acid substitutions for the prevention, delay in the onset of, or early amelioration of type 1 diabetes in a mammal, in particular a human being, at risk of developing said disease.

20

In another aspect, the present invention concerns naturally occurring insulins which are hormonally inactive in mammals, in particular in human beings, for the prevention, delay in the onset of, or early amelioration of type 1 diabetes in a mammal, in particular in a human being.

25

In a further aspect, the present invention relates to a pharmaceutical composition comprising a hormonally inactive insulin.

30

In a still further aspect, the present invention relates to the use of a hormonally inactive insulin for producing a pharmaceutical composition for preventing or delaying the onset of a disease having the characteristics of type 1 diabetes or for ameliorating an early stage thereof in a mammal, in particular a human being, at risk of developing said disease.

35

As used in the present description, the expression "a hormonally active insulin" designates a naturally occurring insulin or an insulin analogue with a biological activity

(hypoglycaemic effect) which makes it potentially useful in the treatment of manifest type 1 diabetes in a mammal, particularly in a human being.

5 As used in the present description, the expression "a hormonally inactive insulin" designates a naturally occurring insulin or an insulin analogue with a biological activity (hypoglycaemic effect) which is so low that it cannot be considered as potentially useful in the treatment
10 of manifest type 1 diabetes in a mammal, particularly in a human being. According to the present invention, such hormonally inactive insulins have an *in vitro* activity which is less than 7%, preferably less than 3%, more preferred less than 1%, still more preferred less than 0.1% of the
15 activity exhibited by human insulin when tested in either the mouse free fat cell assay as described by Vølund *et al.* (1991) *Diabetic Medicine* **8**, 839-847 with reference to Moody *et al.* (1974) *Horm. Metab. Res.* **6**, 12-16 and Vølund (1978) *Biometrics* **34**, 357-365 or in the receptor binding assay as
20 described by Andersen *et al.* (1992) *J. Biol. Chem.* **267**, 13681-13686.

In the present specification, whenever the term insulin is used in a plural or a generic sense it is intended to
25 encompass both naturally occurring insulins and insulin analogues unless otherwise specified. By "insulin analogue" as used herein is meant a polypeptide having a molecular structure similar to that of human insulin including the disulphide bridges between Cys^{A7} and Cys^{B7} and between Cys^{A20}
30 and Cys^{B19} and in most cases also an internal disulphide bridge between Cys^{A6} and Cys^{A11}. Hormonally inactive insulins can formally be derived from human insulin by deleting and/or substituting a moderate number of amino acid residues. In the insulin analogues of the present Table 1
35 the sum of the number of deletions and substitutions is not higher than 7.

In one preferred embodiment, the insulin analogues of the

invention contain no deletions and only two substitutions.

In another preferred embodiment, the insulin analogues of the invention contain no deletions and only one substitution.

Specifically exemplified in the present examples is the use of a human insulin analogue, Asp^{B25} human insulin, to effectively protect against the development of IDD. The hormonally inactive insulins disclosed herein effectively block or immunologically override the autoimmune reaction that causes destruction of the insulin-secreting cells of the pancreas, without exposing the treated individuals to the adverse effects of insulin therapy.

Treatment with compositions of the subject invention obviates the need for daily insulin therapy, the need for repeated blood glucose monitoring, and the restrictions of diet and exercise imposed upon all patients with IDD receiving insulin replacement therapy. Further, and most importantly, the complications of IDD, which are directly attributable to the duration and severity of overt diabetes and hyperglycaemia, will be inhibited. Those who would be likely to benefit most from the treatment would be those with a high genetic risk profile, such as a five-fold or greater risk as determined by HLA and/or other genotyping studies, as well as those found to have a high predictive profile as determined through specific autoantibody analysis, e.g., GAD₆₅ autoantibodies.

BRIEF DESCRIPTION OF THE DRAWINGS

The present invention is further illustrated with reference to the appended drawings wherein

Figure 1 shows the effect on the blood glucose level of daily injections of buffer, various doses of Asp^{B25} human

insulin (code: X38) and various doses of human insulin, respectively;

Figure 2 is a graph showing the effects of subcutaneous administration of human insulin, Asp^{B25} human insulin (code: X38) and buffer on diabetes in NOD mice;

Figure 3 is a graph showing the effects of subcutaneous administration of human insulin, Asp^{B25} human insulin (code: X38) and buffer on survival in NOD mice;

Figure 4 is a graph showing the effects of subcutaneous administration of insulin, Asp^{B25} human insulin (code: X38) and buffer in the initial stages of diabetes in NOD mice; and

Figure 5 is a graph showing the effects of oral administration of insulin or buffer on blood glucose levels in NOD mice.

DETAILED DESCRIPTION OF THE INVENTION

The subject invention concerns insulin compositions uniquely suited for use in new, clinically important, immunomodulating therapies that selectively inhibits beta-cell destructive immune responses without inducing insulin inherent side effects and a state of generalized immunosuppression replete with its serious side effects. The use of the insulin compositions of the subject invention represent an important advance in the development of preventive, interventive or ameliorating therapies for diabetes.

Another critical distinguishing feature of the subject invention is the use of insulin before the appearance of clinical symptoms of IDD. Clearly, because of the metabolic activities of insulin, it is not common practice to

administer insulin unless there is already clinical evidence of pancreatic islet cell destruction and insulin insufficiency. According to the subject invention, however, modified insulin or a fragment thereof, is best administered
5 before any symptoms appear.

Methods of analysing the biological activity (hypoglycaemic effect) of insulins are well known, and include receptor binding assays using whole cells or soluble receptors, the
10 mouse free fat cell assay, euglycaemic pig clamp assay, intravenous rabbit blood glucose assay and the subcutaneous mouse blood glucose assay (Vølund et al. (1991) *Diabetic Medicine*, **8**, 839-847, Brange et al. (1990) *Diabetes Care* **13**, 923-954, Drejer (1992) *Diabetes/Metabolism Reviews* **8**, 259-
15 286).

In a preferred embodiment, the insulin analogues of the invention are selected from the insulin analogues listed in Table 1. The biological activities stated in Table 1 are
20 given relative to human insulin (=1.00) using either receptor binding or lipogenesis assays.

Table 1

25	Insulin analogue	Activity relative to human insulin (= 1.00)
	desA1 human insulin;	0.006
30	des(A1-A2) human insulin;	0.0014
	des(A1-A3) human insulin;	0.0011
	desA21 human insulin;	0.04
	des(B1-B5) human insulin;	0.05
	des(B1-B6) human insulin;	0.0004
35	des(B23-B30) human insulin;	0.001
	des(B24-B30) human insulin;	0.002
	des(B25-B30) human insulin;	0.06
	Gly ^{A2} human insulin;	0.01

	Ala ^{A2} human insulin;	0.006
	Nle ^{A2} human insulin;	0.01
	Thr ^{A2} human insulin;	0.001
	Pro ^{A2} human insulin;	0.01
5	D-allo Ile ^{A2} human insulin;	0.002
	Nva ^{A3} human insulin;	0.014
	Nle ^{A3} human insulin;	0.0038
	Leu ^{A3} human insulin;	0.0018
	Val ^{A2} , Ile ^{A3} human insulin;	0.021
10	Abu ^{A2} , Abu ^{A3} human insulin;	0.0041
	Gly ^{A2} , Gly ^{A3} human insulin;	0.00014
	D-Cys ^{A6} human insulin;	0.0023
	D-Cys ^{A6} , D-Cys ^{A11} human insulin;	0.00027
	Ser ^{A6} , Ser ^{A11} , des(A8-A10) human insulin;	0.0001
15	D-Cys ^{A7} human insulin;	0.002
	D-Cys ^{A11} human insulin;	0.007
	Leu ^{A19} human insulin;	0.001
	Gly ^{B6} human insulin;	0.0005
	Glu ^{B12} human insulin;	0.0004
20	Asn ^{B12} human insulin;	0.002
	Phe ^{B12} human insulin;	0.002
	D-Ala ^{B12} human insulin;	0.007
	Asp ^{B25} human insulin (code: X38).	0.001
25	The insulin analogues of the present invention can be prepared by methods known per se, see for example Märki et al. (1979) Hoppe-Seyler's Z. Physiol. Chem. 360 , 1619-1632; Kitagawa et al. (1984) Biochemistry 23 , 4444-4448; Schwartz et al. (1978) Biochemistry 17 , 4550-4556; Nakagawa et al. (1991) J. Biol. Chem. 266 , 11502-11509; Schwartz et al. (1981) Int. J. Pept. Protein Res. 17 , 243-255; Hu et al. (1993) Biochemistry 32 , 2631-2635; Nakagawa et al. (1986) J. Biol. Chem. 261 , 7332-7341; and Riemen et al. (1983) Biochemistry 22 , 1507-1515.	

35

Pharmaceutical compositions

Pharmaceutical compositions containing an insulin analogue

according to the present invention may be administered orally or parenterally to a person in need of such a treatment. Parenteral administration may be performed by subcutaneous, intramuscular or intravenous injection by means of a syringe, optionally a pen-like syringe. Alternatively, parenteral administration can be performed by means of an infusion pump or a needleless injection device. A further option is to administer a composition which may be a powder or a liquid containing the insulin analogue in the form of a nasal spray. As a still further option, it may also be possible to administer the insulin analogue transdermally.

Pharmaceutical compositions containing an insulin analogue of the present invention may be prepared by conventional techniques, e.g. as described in *Remington's Pharmaceutical Sciences* (1985).

Thus, an injectable composition of an insulin analogue of the invention can be prepared using the conventional techniques of the pharmaceutical industry which involves dissolving and mixing the ingredients as appropriate to give the desired end product.

According to one procedure, the insulin is dissolved in a volume of water which is somewhat below the final volume of the composition to be prepared. An isotonic agent, a preservative, a buffer and optionally other auxiliary agents are added as required and the pH value of the solution is adjusted - if necessary - using an acid, e.g. hydrochloric acid, or a base, e.g. aqueous sodium hydroxide, as needed. Finally, the volume of the solution is adjusted with water to give the desired concentration of the ingredients.

Examples of isotonic agents are sodium chloride, mannitol and glycerol.

Examples of preservatives are phenol, m-cresol, methyl p-hy-

droxybenzoate and benzyl alcohol.

Examples of suitable buffers are sodium acetate, sodium citrate and sodium phosphate. Certain of the above-mentioned
5 auxiliary agents may also be added in order to improve the stability of the composition. Typically, the insulin analogues of the invention are present in the compositions in the form of zinc complexes which may contain from about 0.25 to about 0.75 zinc ions per insulin monomer. Such
10 complexes may further contain phenol and/or m-cresol.

According to one preferred aspect of the present invention, there is provided a pharmaceutical composition of an insulin analogue of the invention in the form of a solution
15 containing hexameric complexes of the said analogue. Typically the hexameric complexes are stabilized by two or more zinc ions and three or more molecules of a phenolic compound like phenol or m-cresol or mixtures thereof per hexamer.

20 A composition for nasal administration of an insulin analogue may, for example, be prepared as described in European Patent No. 272097 (to Novo Nordisk A/S).

25 The insulin analogues of this invention can be used in the prevention of diabetes. The particular insulin analogue to be used and the optimal dose level for any patient will depend on a variety of factors including the efficacy of the specific insulin analogue employed, the age, body weight,
30 physical activity, and diet of the patient and on a possible combination with other drugs. It is recommended that the dosage of the insulin analogue of this invention be determined for each individual patient by those skilled in the art.

35 The present invention is further illustrated by the following examples which, however, are not to be construed as limiting in any way the claimed scope of protection of

the invention. The features disclosed in the foregoing description and in the following examples may, both separately and in any combination thereof, be material for realizing the invention in diverse forms thereof.

5

EXAMPLES

Units of insulin have been converted to mg by using the
10 equation $26 \text{ U} = 1 \text{ mg}$.

EXAMPLE 1

Effect on the blood glucose level of daily administration of
15 Asp^{B25} human insulin to NOD mice.

Groups of 5 female NOD mice (from Bomnice, Bomholtgård, Ry,
20 Denmark), 9 weeks old were treated daily, 7 days a week, for
26 days with subcutaneous injection of either buffer, human
insulin (3.8 mg/kg body weight or 0.6 mg/kg) or Asp^{B25} human
insulin (code: X38) (17.3 mg/kg, 5.8 mg/kg or 0.6 mg/kg). On
25 day seven, an initial blood glucose value was determined at
7:00 am, the injection was given at 7:45 and a further
determination of the blood glucose value was made 6 hours
later. The results are summarised Figure 1 which clearly
demonstrates that even extremely high doses of X38 do not
affect the blood glucose levels. In contrast, insulin (0.6
30 mg/kg) has a marked hypoglycaemic effect. Furthermore, 4 of
the 5 animals receiving insulin (3.8 mg/kg) died of
hypoglycaemia after 3-5 days of treatment and animal number
five had to be sacrificed on day seven due to symptoms of
hypoglycaemia. None of the mice treated with X38, even in
35 the extremely high doses died or displayed hypoglycaemia
during the 26 days observation period.

EXAMPLE 2

Prophylactic effect of daily administration of Asp^{B25} human insulin to NOD mice.

5
—

Female NOD (non-obese diabetic) mice, 4 weeks old, were obtained from Bommice (Bomholtgård, Ry, Denmark). At the beginning of the experiment there were a total of 30 animals in each of the three groups mentioned below. From 5 weeks of age the animals were injected daily for 7 days a week, with either human insulin (Protaphane®, Novo Nordisk A/S, Denmark) 0.77 mg/kg body weight of a 0.15 mg/ml preparation, Asp^{B25} human insulin (code: X38) 0.77 mg/kg body weight of a 0.15 mg/ml preparation or an equal volume of the buffer used for diluting the Protaphane® insulin. The treatment was continued until the mice were 32 weeks old, at which age treatment was stopped and the mice observed until they were 40 weeks old. Beginning when the mice were 10 weeks old, the urine was tested weekly for the presence of glucose using glucosuria test tape (Tes-Tape, Lilly, Indianapolis, IN). If an animal tested positive in the glucose urine test, a blood sample was taken and blood glucose measured using a glucose analyzer. If the blood glucose was 15 mmol/l or higher, the animal was classified as diabetic and then sacrificed.

As shown in Figure 2, a high proportion of the buffer treated animals developed diabetes at an incidence and age at onset similar to that observed in the parent female NOD mouse colony and as reported in the literature. Daily treatment with insulin from 5 weeks of age in doses as previously reported (Atkinson et al. (1990) Diabetes **39**, 933-937, Gotfredsen et al. (1985) Diabetologia **28**, 933-935), significantly reduced the incidence of diabetes ($p < 0.05$) at 31 weeks of age. Termination of treatment was not followed by an abrupt increase in diabetes incidence, and at 40 weeks of age the incidence was still significantly ($p < 0.05$) lower than in the buffer treated animals. Surprisingly, daily

injections of X38 prevented the development of diabetes to the same, or higher degree as insulin did. Thus, at 31 weeks of age after 26 weeks of treatment, the diabetes incidence was highly significantly ($p < 0.01$) lower in the X38 treated animals than in the buffer treated, but not different from that in the insulin treated group. After termination of treatment only 1 case of diabetes occurred, thus at 40 weeks of age the diabetes incidence in the X38 treated animals was still highly significantly lower than in the buffer treated group ($p < 0.01$). At onset of diabetes in these NOD mice, blood glucose values are typically 20 mmol/l or higher. At the end of the experiment, the blood glucose values were determined in non-diabetic animals and all were in the range of 7 - 10 mmol/l (data not shown). Three times during the treatment period, treatment was discontinued for one day and blood glucose values determined 48 hours after the last injection with insulin, X38 or buffer. No previously non-diabetic animals turned out as diabetic following the temporary discontinuation of the treatment. Thus, the treatment was not simply masking hyperglycaemia.

A previous study (Atkinson et al. (1990) Diabetes 39, 933-937) has shown that for prophylactic subcutaneous insulin to be active, high sublethal doses are required. Figure 3 confirms this observation, since 50% of the insulin treated mice died of hypoglycaemia during the treatment period. In contrast, even though X38 provided the same protection against the development of diabetes as insulin did, only one X38 treated animal died. This is not statistically different from the result in the buffer treated group.

These results demonstrate that diabetes can be prevented using a hormonally inactive insulin analogue without the adverse effects of continuous hypoglycaemia due to insulin treatment.

EXAMPLE 3

Effect on diabetes of daily, prophylactic administration of Asp^{B25} human insulin in the initial stage of diabetes in NOD mice.

5

Female NOD (non-obese diabetic) mice, 4-5 weeks old, were obtained from the Jackson Laboratory (Bar Harbour, ME). At the beginning of the experiment, there were a total of 20 animals in each of the three groups mentioned below. From 10 weeks of age, at an age where there are manifest signs of cellular infiltration in the islets of Langerhans and beginning beta cell destruction and by which age blood glucose levels are at hyperglycaemic concentrations in a certain percentage of animals prior to the onset of diabetes, the animals were injected daily for 7 days a week, with either human insulin (Protaphane® Novo Nordisk A/S, Denmark), 0.77 mg/kg body weight of a 0.15 mg/ml preparation, Asp^{B25} human insulin (code: X38), 0.77 mg/kg body weight of a 0.15 mg/ml or an equal volume of the buffer used for diluting the Protaphane® insulin. The treatment was continued until the mice were 37 weeks old, at which age treatment was stopped. The mice were observed until they were 39 weeks old. Beginning when the mice were 10 weeks old, the urine was tested weekly for the presence of glucose using glucosuria test tape (Tes-Tape, Lilly, Indianapolis, IN). If an animal tested positive in the glucose urine test, a blood sample was taken and blood glucose measured using a glucose analyzer. If the blood glucose was 15 mmol/l or higher, the animal was classified as diabetic and sacrificed.

As shown in Figure 4, a high proportion of the buffer treated animals developed diabetes with a frequency similar to that in other female NOD mice from the same colony. Daily treatment with insulin from the age of 10 weeks reduced the frequency of diabetes compared to buffer treated and untreated animals. This effect was demonstrable from the age

of 15 weeks and persisted until the experiment was terminated. Surprisingly, daily injections of X38 reduced the frequency of diabetes to the same, or higher degree as insulin did in the same age interval. After termination of the treatment, no cases of diabetes occurred in the insulin or X38 treated groups of mice. At the onset of diabetes in these NOD mice, blood glucose values were typically 20 mmol/l or higher. At the end of the experiment, the blood glucose values were determined in non-diabetic animals and all were in the range of 7 - 10 mmol/l (data not shown). Three times during the treatment period, treatment was discontinued for one day and blood glucose values determined 48 hours after the last injection of insulin, X38 or buffer. No previously non-diabetic animals turned out as diabetic following the temporary discontinuation of the treatment. Thus, the treatment was not simply masking hyperglycaemia.

In the group of insulin treated mice, 15% died of hypoglycaemia during the treatment period. In contrast, even though X38 provided the same protection against the development of diabetes as insulin did, none of the X38 treated animals died. None of the buffer treated animals died in the observation period.

EXAMPLE 4

Effect of oral administration of Asp^{B25} human insulin on the blood glucose level in NOD mice.

Oral administration of insulin to NOD mice has been reported to prevent diabetes but not to have any effect on the blood glucose level and hence any direct mitogenic effect in the gut milieu (WO 92/06704). Groups of 10 C57/BL mice (from Bomnice, Bomholtgård, Ry, Denmark), 5 weeks old, were fed either buffer, 1 mg of human insulin or 10 mg of human insulin 2 times a week for 5 weeks. After 4 weeks of treatment, the effect on the blood glucose level in non-

fasted animals was analyzed. An initial blood sample was taken in order to determine the control blood glucose level. 15-30 minutes later the animals were fed by gavage and the blood glucose level was determined after further 15, 60 and 180 minutes. Figure 5 clearly shows a profound immediate reduction in the blood glucose level after feeding 1 mg insulin, while at 60 minutes blood glucose level was similar to that of the buffer treated animals. Feeding of a higher dose, 10 mg insulin had an even more profound effect that lasted for the duration of the experiment. These data demonstrate two important points: 1) Oral administration of insulin has a profound hypoglycaemic effect which is not desirable in a preventive treatment and which in itself is a well known risk factor. 2) The fact that sufficient insulin is present in the gut to provoke a hypoglycaemic response demonstrates that a high concentration of intact insulin is present in the gut. Insulin is a well known growth factor and has a direct mitogenic effect. The consequence of having a high concentration of intact insulin in the gut lumen directly exposing gut epithelium to its effect is not known but could potentially be serious.

EXAMPLE 5

A pharmaceutical composition comprising a solution of 600 nmol/ml of Asp^{B25} human insulin, 0.5 zinc ion per insulin monomer, 16 mM m-cresol, 16 mM phenol, 1.6% of glycerol, 23 mM sodium chloride and 7 mM sodium phosphate.

30

1.2 μ mol of Asp^{B25} human insulin was dissolved in water (0.5 ml) by addition of 30 μ l 0.2 M hydrochloric acid to pH 2.5-3.0 and 60 μ l of 0.01 M zinc acetate was added. To the solution was further added 100 μ l of 0.32 M phenol, 200 μ l of 0.16 M m-cresol, 400 μ l of 8% glycerol, 67 μ l of 0.6 M sodium chloride, and pH value of the solution was adjusted to 7.5 by addition of 0.2 M sodium hydroxide. Finally 140 μ l

23

of 0.1 M sodium phosphate (pH 7.5) was added and the volume adjusted to 2 ml with water.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2223
2